


Stereoselective catalytic in-flow synthesis of chiral molecules

Maurizio Benaglia*

¹Università degli Studi di Milano, Dipartimento di Chimica, Via Golgi 19, 20133, Milano, Italy.

Table of contents



Abstract

Continuous-flow systems have emerged as a powerful technology for performing chemical transformations and have recently attracted attention also for the preparation of chiral APIs (active pharmaceutical ingredients).[1]

Recently developed stereoselective transformations performed in chiral organocatalytic reactors under continuous flow conditions will be presented. Both packed-bed and monolithic reactors, either made by silica and/or organic polymers, have been successfully used in metal-free reactions promoted by immobilized enantiopure primary or secondary amines.[2]

The possibility to perform organocatalytic reactions in (micro)-mesoreactors and to synthesize in flow-mode chiral intermediates of pharmaceutical interest will be also discussed.[3]

On the other hand, 3D-printing technology allows chemists to build devices with high precision and well-defined architecture. Stereoselective catalytic in-flow reactions in 3D-printed reactors will be also briefly highlighted.[4]

The combination of these two only partially explored technologies in stereoselective organic synthesis opens new and intriguing possibilities; the fabrication of *ad hoc* designed reactors and other devices, to perform at best different reactions becomes now feasible and gives new impulse to the use of enabling technologies in the synthesis of complex molecules.[5]

Keywords: organocatalysis; flow chemistry; stereoselectivity; chiral pharmaceutical compounds; microreactors.

Acknowledgements

MB thanks Università degli Studi di Milano and Regione Lombardia for a POR-FESR grant.

Reference

- [1] a) B. Gutmann, D. Cantillo, C. O. Kappe, *Angew. Chem. Int. Ed.* **2015**, *54*, 6688-6728; b) R. Porta, M. Benaglia, A. Puglisi, *Org. Process Res. Dev.* **2016**, *20*, 2-25.
 [2] a) R. Porta, M. Benaglia, R. Annunziata, A. Puglisi, G. Celentano, *Adv. Synth. Catal.*, **2017**, 359, DOI: 10.1002/adsc.201700376; b) D. Brenna, R. Porta, E. Massolo, L. Raimondi, M. Benaglia, *ChemCatChem* **2017**, DOI: 10.1002/cctc.201700052; c) R. Porta, M. Benaglia, F. Coccia, F. Cozzi, A. Puglisi, *Adv. Synth. Catal.*, **2015**, *357*, 377-383.

- [3] b) S. Rossi, A. Puglisi, M. Benaglia, D. M. Carminati, D. Intrieri, E. Gallo, *Catal. Sci. Technol.*, **2016**, *6*, 4700-4704d) D. Brenna, M. Benaglia, R. Porta, S. D. Fernandes, A. J. Burke, *Eur. J. Org. Chem.*, **2016**, doi:10.1002/ejoc.201601268.
 [4] a) S. Rossi, D. Brenna, R. Porta, A. Puglisi, M. Benaglia, *Angew. Chem. Int. Ed.* **2017**, *56*, 4290-4294; b) V. Dragone, V. Sans, M. H. Rosnes, P. J. Kitson, L. Cronin, *Beilstein journal of organic chemistry* **2013**, *9*, 951-959
 [5] Review: S. V. Ley, D. E. Fitzpatrick, R. J. Ingham, R. M. Myers, *Angew. Chem. Int. Ed.* **2015**, *54*, 3449.